

# ACMC Intubation Checklist

- **Sterile Cock Pit Announced**
- **Suction set up**
- **Oxygenation** (pre-ox and apneic-ox)
  - NRB at flush rate OR BVM if apneic
  - NC at flush rate
- **Airway Equipment Ready**
  - Laryngoscope
  - ETT + stylet + 10 cc syringe
  - Bougie
  - BVM + peep valve
  - iGel
  - Trach tape/holder
- **Positioning of patient optimized**
  - Ear to sternal notch
  - C-collar off, in-line stabilization held
- **Meds & IV ready**
- **Monitors attached (pulse ox opposite BP cuff)**
- **End tidal CO2 (to Zoll monitor and button on)**

Adult RSI Doses	60 kg (small)	80 kg (med)	100 kg (large)
Ketamine 1.0 mg/kg (10 mg/mL)	60 mg (6 mL)	80 mg (8 mL)	100 mg (10 mL)
Etomidate 0.3 mg/kg (2 mg/mL)	20 mg (10 mL)	25 mg (12.5 mL)	30 mg (15 mL)
Succinylcholine 1.5 mg/kg (20 mg/mL)	90 mg (4.5 mL)	120 mg (6 mL)	150 mg (7.5 mL)
Rocuronium 1.2 mg/kg (10 mg/mL)	70 mg (7 mL)	95 mg (9.5 mL)	120 mg (12 mL)

	Mixing	Dosing
Push dose epi	1 mL code epi in 10 cc flush (10 mcg/mL)	10-20 mcg q2-5 min
Push dose neo	1 mL neo into 100 cc NS (100 mcg/mL)	100-200 mcg q2-5min